Table 6.6g
Immunosuppression Use for Maintenance Between Discharge and One Year Following Transplantation, 1997 to 2006
Recipients with Pancreas Transplant Alone (PTA) Transplants

	Year of Transplant									
	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Transplants	79	88	139	134	167	181	160	183	199	173
Tx with Follow-up Immunosuppression Info	46	63	109	96	122	141	115	137	148	122
Corticosteroids										
Any in Category Steroids	100.0%	100.0% 100.0%	100.0% 100.0%	96.9% 96.9%	92.6% 92.6%	86.5% 86.5%	56.5% 56.5%	56.9% 56.9%	54.1% 54.1%	60.7% 60.7%
Cyclosporine										
Any in Category	23.9%	19.0%	8.3%	8.3%	5.7%	8.5%	7.0%	3.6%	2.0%	2.5%
Sandimmmune	2.2%	0.0%	0.0%	2.1%	0.0%	0.0%	0.0%	0.0%	0.7%	0.8%
Neoral	21.7%	19.0%	8.3%	5.2%	5.7%	6.4%	6.1%	3.6%	1.4%	1.6%
Gengraf	0.0%	0.0%	0.0%	1.0%	0.0%	1.4%	0.9%	0.0%	0.0%	0.0%
Eon	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%	0.0%	0.0%	0.0%	0.0%
Tacrolimus										
Any in Category	80.4%	96.8%	96.3%	95.8%	98.4%	95.0%	84.3%	85.4%	98.0%	97.5%
Tacrolimus	80.4%	96.8%	96.3%	95.8%	98.4%	95.0%	84.3%	85.4%	98.0%	97.5%
Antimetabolites										
Any in Category	91.3%	93.7%	93.6%	81.3%	73.8%	63.8%	60.0%	59.1%	62.8%	59.8%
Mycophenolate Mofetil	78.3%	81.0%	91.7%	76.0%	70.5%	63.8%	59.1%	53.3%	55.4%	51.6%
Mycophenolate Sodium	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	4.1%	6.6%
Azathioprine	13.0%	12.7%	1.8%	5.2%	3.3%	0.0%	0.9%	5.8%	3.4%	1.6%
mTOR Inhibitors										
Any in Category	0.0%	0.0%	17.4%	27.1%	16.4%	33.3%	24.3%	26.3%	27.0%	30.3%
Sirolimus	0.0%	0.0%	17.4%	27.1%	16.4%	33.3%	24.3%	26.3%	27.0%	30.39
Everolimus	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes

Percentages are calculated based on the total number of patients with any immunosuppression use between discharge and one year following transplantation.

Individual drug percentages will not necessarily add up to the drug category percentages, as patients may be prescribed more than one drug within the same category.